

1. **B.** Choice A is odd; choices C and D are even. Choice B is neither as replacing x with $-x$ does not give the original function or the negative of the original function.
2. **B.** $12 = \sqrt{(x-6)^2 + (9+1)^2} = \sqrt{x^2 - 12x + 136} \rightarrow 144 = x^2 - 12x + 136 \rightarrow x^2 - 12x - 8 = 0 \Rightarrow x = 6 \pm 2\sqrt{11}$.
3. **C.** A sketch of each graph shows that they will intersect two times.
4. **A.** Using the first equation, $f(-1) = -1 + 3 = 2$. Now, find where the second function will have a value of 2:
 $2(-1) - c = 2 \Rightarrow c = -4$.
5. **B.** When reflected over $(4, 1)$, the new center will be at $(8, 2)$, yielding the new equation $(x-8)^2 + (y-2)^2 = 25$, which is $x^2 + y^2 - 16x - 4y + 43 = 0$. The ordered quadruple is $(1, -16, -4, 43)$, the sum of whose elements is 24.
6. **B.** Since the negative r -value directs the ray in the opposite direction, $\frac{5\pi}{6}$ is a correct angle.
7. **E.** $2^x = 2^{2x} - 12 \rightarrow 2^{2x} - 2^x - 12 = 0 \rightarrow (2^x - 4)(2^x + 3) = 0 \rightarrow 2^x = 4, 2^x = -3 \Rightarrow x = 2$. When $x = 2, y = 4$, so the sum is 6.
8. **C.** An x -axis reflection changes $f(x)$ to $-f(x)$, and the shift right 7 units changes the latter function to $-f(x-7)$.
9. **A.** The x -intercept is found by $0 = \sqrt{x+9} - 5 \rightarrow 25 = x+9 \rightarrow x = 16$; the y -intercept is found by $y = \sqrt{0+9} - 5 = -2$. Their product is -32 .
10. **D.** The slopes of inverse linear functions are reciprocals of one another. The slope of the given line segment is $\frac{1}{3}$, so the inverse function has slope 3.
11. **A.** To obtain the top part of the ellipse, solve for y and take only the positive square root:
 $\frac{(y-3)^2}{9} = 1 - \frac{(x+2)^2}{4} \rightarrow \frac{(y-3)}{3} = \sqrt{\frac{4-(x+2)^2}{4}} \Rightarrow y = \frac{3}{2}\sqrt{-x^2 - 4x + 3}$.
12. **C.** The asymptotes are found by using the “parentetical” part of the equation and the square root of the denominators: $(y+2) = \pm \frac{\sqrt{4}}{\sqrt{2}}(x-1) \Rightarrow (y+2) = \pm\sqrt{2}(x-1)$.
13. **C.** The first equation has slope $-\frac{1}{3}$ and y -intercept $\frac{2}{3}$. The second equation has slope $\frac{-2}{a^2+2}$ and y -intercept $\frac{a+2}{a^2+2}$. To be parallel, the slopes must be equal but the y -intercepts cannot be equal. Using the slopes and solving for a : $-\frac{1}{3} = \frac{-2}{a^2+2} \rightarrow a^2 = 4 \rightarrow a = \pm 2$. Using the y -intercepts and solving for a , $\frac{2}{3} = \frac{a+2}{a^2+2} \rightarrow 2a^2 - 3a - 2 = 0 \rightarrow (2a+1)(a-2) = 0 \rightarrow a = -\frac{1}{2}, 2$. To be parallel, a must be -2 , and to be the same line, a must be 2. Their sum is 0.

14. **A.** Multiply the second equation by -2 and use elimination. This yields $\frac{19}{y} = -2 \rightarrow y = -\frac{19}{2}$. Substituting to find x , we get $\frac{4}{x} - \frac{14}{19} = 10 \rightarrow \frac{4}{x} = \frac{204}{19} \rightarrow x = \frac{19}{51}$. The sum is $\frac{38}{102} - \frac{969}{102} = -\frac{931}{102}$.
15. **E.** The slope of the radius from the center to the point of tangency is $-\frac{3}{4}$ so the slope of the tangent line is $\frac{4}{3}$. Using point-slope form we have $y - 3 = \frac{4}{3}(x + 4) \rightarrow 3y - 9 = 4x + 16 \Rightarrow 4x - 3y = -25$.
16. **D.** If the y -intercept is $(0, h)$, then the area of the triangle is $T = \frac{1}{2}ah$, so $h = \frac{2T}{a}$. The slope of the line is $\frac{h}{a}$, and $\frac{h}{a} = \frac{2T}{a^2}$. The slope-intercept form of the line is $y = \frac{2T}{a^2}x + \frac{2T}{a} \rightarrow 2Tx - a^2y + 2aT = 0$.
17. **A.** Substituting the given points for (x, y) , we obtain $\begin{cases} P + Q = -4 \\ P - Q = 2 \end{cases}$. By elimination we have $2P = -2 \rightarrow P = -1$. By substitution we have $-1 + Q = -4 \rightarrow Q = -3$. $P + Q = -1 + (-3) = -4$.
18. **C.** Take the determinant of the matrix formed by the ordered triples:
By the diagonal method, $\begin{vmatrix} 3 & 2 & 1 \\ -1 & 3 & 0 \\ 2 & 2 & 5 \end{vmatrix} = (45 + 0 - 2) - (6 + 0 - 10) = 47$.
19. **A.** $f(x) = \frac{x^3 + 7x^2 + 16x + 12}{x^2 + 6x + 8} = \frac{(x+2)^2(x+3)}{(x+2)(x+4)}$. There is one vertical asymptote at $x = -4$. Since an $(x+2)$ factor cancels out, there is a hole in the graph at $(-2, 0)$. The highest power of x is in the numerator so there is no horizontal asymptote.
20. **C.** For the x -coordinate, $|12 - 4| = 8$. Five-sixths of 8 is $\frac{20}{3}$. $\frac{20}{3} + 4 = \frac{32}{3}$. For the y -coordinate, $|-6 - 18| = 24$. Five-sixths of 24 is 20. $18 - 20 = -2$. (The y -coordinate becomes more negative so we have to subtract the 20.)
21. **D.** Notice the capital S, signifying the use of principal values.
22. **E.** Using elimination we have $x^2 - x = 0 \rightarrow x(x-1) = 0 \Rightarrow (0, 10), (1, 9)$. The distance between the points is $\sqrt{(1-0)^2 + (9-10)^2} = \sqrt{2}$.
23. **B.** Any point on the angle bisector is equidistant from the rays (lines) that form the angle. Using the distance from a point to a line formula gives us $\frac{3x-4y+2}{5} = \pm \frac{5x+12y-3}{13}$. From here we have two equations:
 $39x - 52y + 26 = 25x + 60y - 15 \Rightarrow 14x - 112y + 41 = 0$ and $-39x + 52y - 26 = 25x + 60y - 15 \Rightarrow 64x + 8y + 11 = 0$.
24. **D.** S has polar coordinates $(4, 0^\circ)$; U has coordinates (r, θ) , $r \geq 0$. We need $d(0, U) + d(U, S) = 6 \rightarrow$


$$r + \sqrt{4^2 + r^2 - 2(4)r \cos(0 - \theta)} = 6 \rightarrow 16 + r^2 - 8r \cos \theta = 36 - 12r + r^2 \rightarrow 3r - 2r \cos \theta = 5 \Rightarrow r = \frac{5}{3 - 2 \cos \theta}.$$

25. **B.** Using (1, 2, 4) as our “first” point, we must subtract the first point’s coordinates from the second point’s coordinates to get the distance: $(4 - 1, 2 - 2, -1 - 4) = (3, 0, -5)$. Although we could switch the first and second points, only one correct answer is given: $\mathbf{r} = (1 + 3t)\mathbf{i} + 2\mathbf{j} + (4 - 5t)\mathbf{k}$. The other answer would have been $\mathbf{r} = (4 - 3t)\mathbf{i} + 2\mathbf{j} + (-1 + 5t)\mathbf{k}$

26. **D.** Each of the ten central angles has measure 36° , and each isosceles triangle has area $\frac{1}{2}(4)^2 \sin 36^\circ$. Since there are ten triangles that form the hexagon, the area is $10\left(\frac{1}{2}(4)^2 \sin 36^\circ\right) = 80 \sin 36^\circ$.

27. **C.** To eliminate the xy term, the angle can be found by $\frac{1}{2} \arctan \frac{B}{A-C} \rightarrow \frac{1}{2} \arctan \frac{\sqrt{3}}{2-3} \rightarrow \frac{1}{2}(120^\circ) \Rightarrow 60^\circ$.


28. **E.** If R is the radius of the circle, then the circumference is $2\pi R = 3 + 4 + 5 = 12$, so $R = \frac{6}{\pi}$. The arcs make up $\frac{1}{4}, \frac{1}{3}$, and $\frac{5}{12}$ of the circle, so their respective central angles are $\frac{\pi}{2}, \frac{2\pi}{3}$, and $\frac{5\pi}{6}$ radians. Since the triangles are all isosceles, we can use the formula $\frac{1}{2}R^2 \sin \theta$ for the area of each triangle:



$$\frac{1}{2} \left(\frac{6}{\pi}\right)^2 \left(\sin \frac{\pi}{2} + \sin \frac{2\pi}{3} + \sin \frac{5\pi}{6}\right) = \frac{18}{\pi^2} \left(1 + \frac{\sqrt{3}}{2} + \frac{1}{2}\right) = \frac{9}{\pi^2} (3 + \sqrt{3}).$$


29. **D.** There are 16 lattice points in the given region, and there are ${}_{16}C_3 = 560$ sets of three points. There are four vertical line segments containing four lattice points and four horizontal line segments containing four lattice points, so there are $8\binom{4}{3} = 32$ sets of collinear points that must be subtracted. There are also oblique line segments to consider. Three of them have slope 1 and three have slope -1 .

For each slope there are $\binom{3}{3} + \binom{4}{3} + \binom{3}{3} = 6$ sets of collinear points. Therefore, there are $560 - 32 - 12 = 516$ triangles that satisfy the given condition.



30. **C.** $f(x) = \sqrt{8x - x^2} - \sqrt{14x - x^2 - 48} = \sqrt{16 - (x - 4)^2} - \sqrt{1 - (x - 7)^2}$.

The first radical is the formula for the y -coordinate of the upper half of a circle with center at (4, 0) and radius 4; the second radical is the formula for the y -coordinate of the upper half of a circle with center (7, 0) and radius 1. $f(x)$ is the difference in the heights of the two semicircles. Graphing these two semicircles shows that the function has domain $[6, 8]$ and that the greatest difference must occur at $x = 6$.



$$f(6) = \sqrt{16 - (6 - 4)^2} = \sqrt{12} = 2\sqrt{3}.$$