

#0 Mu CIPHERING

MAΘ National Convention 2011

Find the equation of the tangent line to the graph of $y = \sin x + x$ at the point $(0,0)$.

#0 Mu CIPHERING

MAΘ National Convention 2011

Find the equation of the tangent line to the graph of $y = \sin x + x$ at the point $(0,0)$.

#0 Mu CIPHERING

MAΘ National Convention 2011

Find the equation of the tangent line to the graph of $y = \sin x + x$ at the point $(0,0)$.

#0 Mu CIPHERING

MAΘ National Convention 2011

Find the equation of the tangent line to the graph of $y = \sin x + x$ at the point $(0,0)$.

#1 Mu Ciphering
MA $\text{\textcircled{C}}$ National Convention 2011

Find the range of the function $y = \frac{x+1}{x^2+1}$.

#1 Mu Ciphering
MA $\text{\textcircled{C}}$ National Convention 2011

Find the range of the function $y = \frac{x+1}{x^2+1}$.

#1 Mu Ciphering
MA $\text{\textcircled{C}}$ National Convention 2011

Find the range of the function $y = \frac{x+1}{x^2+1}$.

#1 Mu Ciphering
MA $\text{\textcircled{C}}$ National Convention 2011

Find the range of the function $y = \frac{x+1}{x^2+1}$.

#2 Mu Ciphering**MA \odot National Convention 2011**

Find the slope of the line tangent to the graph

of $xy^2 + \sin x + y = \frac{x}{y^2} + 1$ at the point $(0,1)$.

#2 Mu Ciphering**MA \odot National Convention 2011**

Find the slope of the line tangent to the graph

of $xy^2 + \sin x + y = \frac{x}{y^2} + 1$ at the point $(0,1)$.

#2 Mu Ciphering**MA \odot National Convention 2011**

Find the slope of the line tangent to the graph

of $xy^2 + \sin x + y = \frac{x}{y^2} + 1$ at the point $(0,1)$.

#2 Mu Ciphering**MA \odot National Convention 2011**

Find the slope of the line tangent to the graph

of $xy^2 + \sin x + y = \frac{x}{y^2} + 1$ at the point $(0,1)$.

#3 Mu Ciphering
MA \odot National Convention 2011

Evaluate the integral $\int_0^3 (9 - x^2)^{1/2} dx$.

#3 Mu Ciphering
MA \odot National Convention 2011

Evaluate the integral $\int_0^3 (9 - x^2)^{1/2} dx$.

#3 Mu Ciphering
MA \odot National Convention 2011

Evaluate the integral $\int_0^3 (9 - x^2)^{1/2} dx$.

#3 Mu Ciphering
MA \odot National Convention 2011

Evaluate the integral $\int_0^3 (9 - x^2)^{1/2} dx$.

#4 Mu Cipheryng
MA⊗ National Convention 2011

Find the value of $\lim_{n \rightarrow \infty} \sum_{x=1}^n \left(\frac{x^2 + 2n^2}{n^3} \right)$.

#4 Mu Cipheryng
MA⊗ National Convention 2011

Find the value of $\lim_{n \rightarrow \infty} \sum_{x=1}^n \left(\frac{x^2 + 2n^2}{n^3} \right)$.

#4 Mu Cipheryng
MA⊗ National Convention 2011

Find the value of $\lim_{n \rightarrow \infty} \sum_{x=1}^n \left(\frac{x^2 + 2n^2}{n^3} \right)$.

#4 Mu Cipheryng
MA⊗ National Convention 2011

Find the value of $\lim_{n \rightarrow \infty} \sum_{x=1}^n \left(\frac{x^2 + 2n^2}{n^3} \right)$.

#5 Mu Ciphering

MA \odot National Convention 2011

A man 6 feet tall walks at a rate of 5 ft/s away from a light that is 15 ft above the ground. When he is 10 ft from the base of the light, at what rate, in ft/s, is the length of his shadow changing?

#5 Mu Ciphering

MA \odot National Convention 2011

A man 6 feet tall walks at a rate of 5 ft/s away from a light that is 15 ft above the ground. When he is 10 ft from the base of the light, at what rate, in ft/s, is the length of his shadow changing?

#5 Mu Ciphering

MA \odot National Convention 2011

A man 6 feet tall walks at a rate of 5 ft/s away from a light that is 15 ft above the ground. When he is 10 ft from the base of the light, at what rate, in ft/s, is the length of his shadow changing?

#5 Mu Ciphering

MA \odot National Convention 2011

A man 6 feet tall walks at a rate of 5 ft/s away from a light that is 15 ft above the ground. When he is 10 ft from the base of the light, at what rate, in ft/s, is the length of his shadow changing?

#6 Mu Ciphering**MA@ National Convention 2011**

Let $f(x) = 2x^3 + 6x - 34$ and $g(x) = f^{-1}(x)$.

Find the value of $(g'(-6))^{-1}$.

#6 Mu Ciphering**MA@ National Convention 2011**

Let $f(x) = 2x^3 + 6x - 34$ and $g(x) = f^{-1}(x)$.

Find the value of $(g'(-6))^{-1}$.

#6 Mu Ciphering**MA@ National Convention 2011**

Let $f(x) = 2x^3 + 6x - 34$ and $g(x) = f^{-1}(x)$.

Find the value of $(g'(-6))^{-1}$.

#6 Mu Ciphering**MA@ National Convention 2011**

Let $f(x) = 2x^3 + 6x - 34$ and $g(x) = f^{-1}(x)$.

Find the value of $(g'(-6))^{-1}$.

#7 Mu Ciphering**MA \odot National Convention 2011**

Find the length of the curve $y = \ln|\cos x|$ from
 $x = 0$ to $x = \pi/4$.

#7 Mu Ciphering**MA \odot National Convention 2011**

Find the length of the curve $y = \ln|\cos x|$ from
 $x = 0$ to $x = \pi/4$.

#7 Mu Ciphering**MA \odot National Convention 2011**

Find the length of the curve $y = \ln|\cos x|$ from
 $x = 0$ to $x = \pi/4$.

#7 Mu Ciphering**MA \odot National Convention 2011**

Find the length of the curve $y = \ln|\cos x|$ from
 $x = 0$ to $x = \pi/4$.

#8 Mu Ciphering**MA $\text{\textcircled{C}}$ National Convention 2011**

A rectangle is inscribed in a triangle whose sides have lengths of 6, 8, and 10. Two vertices of the rectangle are on the longest side of the triangle, and one vertex of the rectangle is on each of the other two sides. Find the maximum area of a rectangle that can be inscribed in this manner.

#8 Mu Ciphering**MA $\text{\textcircled{C}}$ National Convention 2011**

A rectangle is inscribed in a triangle whose sides have lengths of 6, 8, and 10. Two vertices of the rectangle are on the longest side of the triangle, and one vertex of the rectangle is on each of the other two sides. Find the maximum area of a rectangle that can be inscribed in this manner.

#8 Mu Ciphering**MA $\text{\textcircled{C}}$ National Convention 2011**

A rectangle is inscribed in a triangle whose sides have lengths of 6, 8, and 10. Two vertices of the rectangle are on the longest side of the triangle, and one vertex of the rectangle is on each of the other two sides. Find the maximum area of a rectangle that can be inscribed in this manner.

#8 Mu Ciphering**MA $\text{\textcircled{C}}$ National Convention 2011**

A rectangle is inscribed in a triangle whose sides have lengths of 6, 8, and 10. Two vertices of the rectangle are on the longest side of the triangle, and one vertex of the rectangle is on each of the other two sides. Find the maximum area of a rectangle that can be inscribed in this manner.

#9 Mu Ciphering

MA Θ National Convention 2011

A trough is 15 ft long and 4 ft across the top. Its ends are isosceles triangles with height 3 ft. Water runs into the trough at a rate of 2.5 cubic ft/min. How fast is the water level rising, in ft/min, when the water is 2 ft deep?

#9 Mu Ciphering

MA Θ National Convention 2011

A trough is 15 ft long and 4 ft across the top. Its ends are isosceles triangles with height 3 ft. Water runs into the trough at a rate of 2.5 cubic ft/min. How fast is the water level rising, in ft/min, when the water is 2 ft deep?

#9 Mu Ciphering

MA Θ National Convention 2011

A trough is 15 ft long and 4 ft across the top. Its ends are isosceles triangles with height 3 ft. Water runs into the trough at a rate of 2.5 cubic ft/min. How fast is the water level rising, in ft/min, when the water is 2 ft deep?

#9 Mu Ciphering

MA Θ National Convention 2011

A trough is 15 ft long and 4 ft across the top. Its ends are isosceles triangles with height 3 ft. Water runs into the trough at a rate of 2.5 cubic ft/min. How fast is the water level rising, in ft/min, when the water is 2 ft deep?

#10 Mu Cipheryng
MA© National Convention 2011

Let S be the region in the first quadrant enclosed by the y -axis and the graphs of $y = \sin x$ and $y = \cos x$. Find the volume of the solid whose base is S and whose cross-sections cut by planes perpendicular to the x -axis are squares.

#10 Mu Cipheryng
MA© National Convention 2011

Let S be the region in the first quadrant enclosed by the y -axis and the graphs of $y = \sin x$ and $y = \cos x$. Find the volume of the solid whose base is S and whose cross-sections cut by planes perpendicular to the x -axis are squares.

#10 Mu Cipheryng
MA© National Convention 2011

Let S be the region in the first quadrant enclosed by the y -axis and the graphs of $y = \sin x$ and $y = \cos x$. Find the volume of the solid whose base is S and whose cross-sections cut by planes perpendicular to the x -axis are squares.

#10 Mu Cipheryng
MA© National Convention 2011

Let S be the region in the first quadrant enclosed by the y -axis and the graphs of $y = \sin x$ and $y = \cos x$. Find the volume of the solid whose base is S and whose cross-sections cut by planes perpendicular to the x -axis are squares.

#11 Mu Ciphering

MA \odot National Convention 2011

Find the area inside the outer loop and outside the inner loop of the polar graph $r = 1 + 2\cos\theta$.

#11 Mu Ciphering

MA \odot National Convention 2011

Find the area inside the outer loop and outside the inner loop of the polar graph $r = 1 + 2\cos\theta$.

#11 Mu Ciphering

MA \odot National Convention 2011

Find the area inside the outer loop and outside the inner loop of the polar graph $r = 1 + 2\cos\theta$.

#11 Mu Ciphering

MA \odot National Convention 2011

Find the area inside the outer loop and outside the inner loop of the polar graph $r = 1 + 2\cos\theta$.

#12 Mu Ciphering
MA@ National Convention 2011

Find the value of $\lim_{x \rightarrow 0} (1 + 2x)^{\csc x}$.

#12 Mu Ciphering
MA@ National Convention 2011

Find the value of $\lim_{x \rightarrow 0} (1 + 2x)^{\csc x}$.

#12 Mu Ciphering
MA@ National Convention 2011

Find the value of $\lim_{x \rightarrow 0} (1 + 2x)^{\csc x}$.

#12 Mu Ciphering
MA@ National Convention 2011

Find the value of $\lim_{x \rightarrow 0} (1 + 2x)^{\csc x}$.

#13 Mu Cipheryng
MA© National Convention 2011

Find the indefinite integral $\int \frac{xe^{2x}}{(2x+1)^2} dx$.

#13 Mu Cipheryng
MA© National Convention 2011

Find the indefinite integral $\int \frac{xe^{2x}}{(2x+1)^2} dx$.

#13 Mu Cipheryng
MA© National Convention 2011

Find the indefinite integral $\int \frac{xe^{2x}}{(2x+1)^2} dx$.

#13 Mu Cipheryng
MA© National Convention 2011

Find the indefinite integral $\int \frac{xe^{2x}}{(2x+1)^2} dx$.

#14 Mu Ciphering**MA© National Convention 2011**

The position of a particle moving on the x -axis is given by $x(t) = \sin t \cos t$ for $t \geq 0$. At the first instant when the acceleration is equal to 1, what is the particle's velocity?

#14 Mu Ciphering**MA© National Convention 2011**

The position of a particle moving on the x -axis is given by $x(t) = \sin t \cos t$ for $t \geq 0$. At the first instant when the acceleration is equal to 1, what is the particle's velocity?

#14 Mu Ciphering**MA© National Convention 2011**

The position of a particle moving on the x -axis is given by $x(t) = \sin t \cos t$ for $t \geq 0$. At the first instant when the acceleration is equal to 1, what is the particle's velocity?

#14 Mu Ciphering**MA© National Convention 2011**

The position of a particle moving on the x -axis is given by $x(t) = \sin t \cos t$ for $t \geq 0$. At the first instant when the acceleration is equal to 1, what is the particle's velocity?

